

Gallbladder Stone



■ Is Gallbladder Stone Dangerous?

Gallstone, if symptomatic, causes acute attack of biliary colic/cholecystitis. Sometimes their migration from gallbladder may lead to blockage of common bile duct or pancreatic duct resulting in serious complications e.g, jaundice, pancreatitis.

■ What is Asymptomatic Gallstone?

It is defined as the detection of gallstones without related symptoms or sequelae such as colic, cholecystitis, cholangitis, or pancreatitis.

■ Is Gallbladder Surgery safe?

Laparoscopic gallbladder surgery is safe and effective. Surgery gets rid of gallbladder together with the stones.

■ Can you live without Gallbladder?

Obviously, when you're living without a gallbladder, it can no longer store bile, as designed. Bile is a liquid that helps you digest fatty foods. ... but most animal species in the world don't have gallbladders. And your gallbladder is an organ you can live without.

■ How does removal of Gallbladder affect a person?

When the gallbladder is removed, bile made by the liver can no longer be stored between meals. Instead, the bile flows directly into the intestine anytime the liver produces it. Thus, there still is bile in the intestine to mix with food and fat. ... The only clear side effect of removal of the gallbladder is diarrhoea.

■ Can you eat normal after Gallbladder Surgery?

Most people don't need to follow a special diet after having surgery to remove their gallbladder, as the gallbladder isn't essential for digestion. You can usually start eating normally a few hours after your operation, although you'll probably prefer to eat small meals to start with.